



PROPICONAZOLE	GROUP	3	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE

SATORI FIT™ FUNGICIDE provides two different modes of action (preventive and curative) delivered systemically via xylem mobility resulting in broad-spectrum, long-lasting residual control of major foliar diseases. It increases yield potential by extending the photosynthesis period for more plant growth.

FEATURES

- Active Ingredients
 - Propiconazole: 1.02 lbs/gal
 - Azoxystrobin: 1.18 lbs/gal
- Formulaton Type: Suspo-Emulsion (SE) Concentrate
- Pack Size: 2x2.5 gal
- Signal Word: Caution

GROWER BENEFITS

- Dual mode of action fungicide for resistance management
 - Quinone Outside Inhibitor (QoI) - Strobilurin
 - DeMethylation Inhibitor (DMI) - Triazole
- Delivers both preventative and curative disease control
- Labeled for use on multiple crops
- Broad-spectrum foliar disease control

KEY CROPS & RATE

Corn: 10.5-14 fl. oz./A Wheat: 7-14 fl. oz./A
 Soybean: 10.5-21 fl. oz./A Rice: 14-27 fl. oz./A

MAXIMIZE PERFORMANCE WITH*



* Confirm adjuvant use is recommended for the intended crop and crop stage prior to use.

KEY DISEASE CONTROL & SUPPRESSION†

KEY DISEASE CONTROL & SUPPRESSION ACROSS CROPS	
CORN	
Gray Leaf Spot	Anthracnose Leaf Blight
Northern Corn Leaf Blight	Tar Spot (2ee)
Rust, Common & Southern	
WHEAT	
Glume Blotch	Tan Spot
Leaf Blight	Rust
Powdery Mildew	
SOYBEAN	
Frogeye Leaf Spot	Cercospora Blight & Leaf Spot
RICE	
Brown Leaf Spot	Sheath Blight
Kernel Smut	Sheath Spot
Leaf Blast	Stem Rot
Narrow Brown Leaf Spot	

† Reference label or applicable 2(ee) for specifics.

