



Dyna-Shield® Imidacloprid 5

Seed Treatment:

This seed has been treated with Imidacloprid:

1-[(6-chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine insecticide at the rate specified by the manufacturer.

Caution: TREATED SEED. DO NOT USE FOR FEED, FOOD OR OIL PURPOSES. STORE AWAY FROM FEED AND FOODSTUFFS. DO NOT USE BAG OR CONTAINER FOR REFILLING WITH FOODSTUFF. KEEP OUT OF REACH OF CHILDREN.

- Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- In the event of a crop failure or harvest of a crop grown from Dyna-Shield Imidacloprid 5 treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, curcubits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat and rice, the minimum plant-back interval is 30 days from the date Dyna-Shield Imidacloprid 5 treated seed was planted. Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.
- The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 pound.
- This seed has been treated with pounds imidacloprid per pounds of seed [or milligrams imidacloprid per seed].
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- Rape greens grown and harvested from Dyna-Shield Imidacloprid 5 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from Dyna-Shield Imidacloprid 5 treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.



Loveland Products, Inc., PO Box 1286, Greeley CO 80632

Dyna-Shield and the Dyna-Shield logo are registered trademarks of Loveland Products, Inc. 070108 V3D 10G09 [K0113LTRPDF]



Dyna-Shield® Imidacloprid 5

Seed Treatment:

This seed has been treated with Imidacloprid:

1-[(6-chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine insecticide at the rate specified by the manufacturer.

Caution: TREATED SEED. DO NOT USE FOR FEED, FOOD OR OIL PURPOSES. STORE AWAY FROM FEED AND FOODSTUFFS. DO NOT USE BAG OR CONTAINER FOR REFILLING WITH FOODSTUFF. KEEP OUT OF REACH OF CHILDREN.

- Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- In the event of a crop failure or harvest of a crop grown from Dyna-Shield Imidacloprid 5 treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, curcubits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat and rice, the minimum plant-back interval is 30 days from the date Dyna-Shield Imidacloprid 5 treated seed was planted. Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.
- The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 pound.
- This seed has been treated with pounds imidacloprid per pounds of seed [or milligrams imidacloprid per seed].
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- Rape greens grown and harvested from Dyna-Shield Imidacloprid 5 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from Dyna-Shield Imidacloprid 5 treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.



Loveland Products, Inc., PO Box 1286, Greeley CO 80632

Dyna-Shield and the Dyna-Shield logo are registered trademarks of Loveland Products, Inc. 070108 V3D 10G09 [K0113LTRPDF]